3-Algebraic structures of the quantum Calogero-Moser model

Chun-Hong Zhang[a,1], Lu Ding[b,2], Zhao-Wen Yan[c,3],
Ke Wu[a,d,4] and Wei-Zhong Zhao[a,d, 5]

[a] *School of Mathematical Sciences, Capital Normal University, Beijing 100048, China*

[b] *Institute of Applied Mathematics, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China*

[c] *School of Mathematical Sciences, Inner Mongolia University, Hohhot 010021, China*

[d] *Beijing Center for Mathematics and Information Interdisciplinary Sciences, Beijing 100048, China*

arXiv:1409.3344v2 [hep-th] 23 Apr 2015

Abstract

We investigate the quantum Calogero-Moser model and reveal its hidden symmetries, i.e., the $W_{1+\infty}$ and Virasoro-Witt 3-algebras. In the large N limit, we note that these two infinite dimensional 3-algebras reduce to the w_∞ and special Virasoro-Witt 3-algebras which satisfy the fundamental identity condition, respectively.

KEYWORDS: Conformal and W Symmetry, Integrable Equations in Physics

[1] zchsxwl@126.com
[2] dinglu@amss.ac.cn
[3] yzw428@126.com
[4] wuke@cnu.edu.cn
[5] Corresponding author: zhaowz@cnu.edu.cn

1.Introduction

The infinite-dimensional algebras play a very important role in the study of integrable systems. The intrinsic connection between the Virasoro algebra and the Korteweg-de Vries (KdV) equation was firstly pointed out by Gervais and Neveu [1, 2]. Then it was shown that the W_N algebra is associated to the generalized KdV hierarchies [3, 4]. It is well-known that all the (generalized) KdV hierarchies can be incorporated into the KP hierarchy. One found that the $W_{1+\infty}$ algebra is intrinsically related to the first Hamiltonian structure of the KP hierarchy [5, 6]. Moreover the second Hamiltonian structure of the KP hierarchy was shown to be isomorphic to $\hat{W}_\infty$ [7], a centerless deformation of W_∞. Calogero-Moser-Sutherland (CMS) models [8, 9, 10] are the important classical and quantum integrable systems in physics, they have attracted a lot of interesting due to their relevance in various fields, such as quantum Hall effect [11, 12, 13, 14], fractional statistics [15, 16], quantum chaos and matrix models [17], black hole [18], Seiberg-Witten theory [19] and string theory [20]. The quantum Calogero-Moser (CM) model is a one-dimensional quantum many-body system with inverse-square long-range interaction. The algebraic structure of this integrable system have been well investigated. The $U(1)$-current [21] and $W_{1+\infty}$ [22] algebraic structures have been found. The $SU(\nu)$ Calogero spin system can be regarded as the generalization of the quantum CM model. Hikami and Wadati [23] found that there is the $su(\nu)$ loop algebraic structure for the $SU(\nu)$ Calogero spin system. Moreover they pointed out that this system also has a higher symmetry, i.e., the $W_{1+\infty}$ algebra. This W algebraic structure unifies the Calogero type and the Sutherland type (trigonometric potential).

Recently 3-algebras have been paid great attention due to a world-volume description of multiple M2-branes proposed by Bagger and Lambert [24], and Gustavsson [25] (BLG). Not as the case of the finite-dimensional 3-algebras, the study of the infinite-dimensional 3-algebras did only make remarkable progress in recent years. The Virasoro-Witt (V-W) algebra is an important infinite-dimensional algebra which is a centerless Virasoro algebra. One found that its 3-algebra is null [26]. Recently Ding et al. [27] investigated the q-deformation of this infinite-dimensional null 3-algebra and constructed a nontrivial q-deformed V-W 3-algebra. A nontrivial V-W 3-algebra was presented through the use of su(1,1) enveloping algebra techniques [28]. In this ternary algebraic structure, there exists a parameter such that it becomes a Nambu 3-algebra only for the special values. Chakrabortty et al. [29] investigated the W_∞ 3-algebra. Although the Filippov condition or fundamental identity (FI) condition [30, 31] was confirmed to fail for this ternary algebra, by applying a double scaling limits on the generators, they derived a w_∞ 3-algebra which satisfies the FI condition. The super w_∞ 3-algebra was constructed in Ref.[30]. It was shown that this super infinite-dimensional 3-algebra satisfies the generalized FI condition. Furthermore a super Nambu-Poisson bracket was proposed there. In terms of this super Nambu-Poisson bracket, a realization of this super w_∞ 3-algebra was also presented.

Besides the infinite-dimensional 3-algebras mentioned above, the Kac-Moody 3-algebra has also been constructed [33].

Quite recently, attempt has been made to establish the connection between the infinite-dimensional 3-algebras and the integrable systems [34, 35]. Chen er al. [34] investigated the classical Heisenberg and w_∞ 3-algebras and established the connection with the dispersionless KdV hierarchy. The bi-Nambu-Hamiltonian structure of the dispersionless KdV system was found. Since the Nambu-Poisson evolution equation involves two Hamiltonians, the deep relationship between these Hamiltonians has also been revealed. Although the progress has been made towards understanding the infinite-dimensional 3-algebras in recent years, to our best knowledge, much less is still known about their connections with the integrable systems. In this paper, we reinvestigate the quantum CM model. By means of the boost and Lax operators, we show that this integrable quantum many-body system has the infinite-dimensional 3-algebraic structures.

2. Quantum Calogero-Moser model

In this section, we shall recall the quantum CM model that will be useful in what follows. For a more detailed description we refer the reader to Ref.[21, 22].

The quantum CM model is an important integrable system in physics. In this integrable system, all particles move on a line interacting through the inverse square interactions. The corresponding Hamiltonian is

$$H = \frac{1}{2}\sum_{j=1}^{N} p_j^2 + \sum_{1\leq j<k\leq N} \frac{a^2-a}{(x_j-x_k)^2}, \tag{1}$$

where $p_j = -\mathbf{i}\frac{\partial}{\partial x_j}$ denotes the momentum operator of the j-th particle, the constants N and a are the particle number and the coupling parameter, respectively. For convenience, we set $\hbar = 1$ and the particle mass $\mathbf{m} = 1$ throughout this paper.

The integrability of the quantum CM model is guaranteed by the existence of a Lax pair. It consists of two $N \times N$ operator-valued matrices L and M,

$$\begin{aligned} L_{jk} &= \delta_{jk}p_j + (1-\delta_{jk})\mathbf{i}a\frac{1}{x_j-x_k}, \\ M_{jk} &= \delta_{jk}a\sum_{l\neq j}\frac{1}{(x_j-x_l)^2} - (1-\delta_{jk})a\frac{1}{(x_j-x_k)^2}. \end{aligned} \tag{2}$$

The operator M satisfies the sum-to-zero condition,

$$\sum_j M_{jk} = \sum_k M_{jk} = 0. \tag{3}$$

For the Lax operators L and M, they satisfy the Lax equation

$$\frac{d}{dt}L_{jk} = \mathbf{i}[H, L_{jk}] = \mathbf{i}[L, M_{jk}]. \tag{4}$$

For the Hamiltonian (1), the corresponding Heisenberg equations of motion are

$$\begin{aligned} \frac{dx_j}{dt} &= \mathbf{i}[H, x_j] = p_j, \\ \frac{dp_j}{dt} &= \mathbf{i}[H, p_j] = \sum_{k\neq j} \frac{2(a^2-a)}{(x_j - x_k)^3}. \end{aligned} \tag{5}$$

By means of the Lax equation (4) and the sum-to-zero condition (3), it is not hard to verify that

$$[H, \sum_{j,k}(L^n)_{jk}] = \sum_{j,k}[L^n, M]_{jk} = 0. \tag{6}$$

From (6), we have the conserved operators of the quantum Calogero-Moser system,

$$I_n = \frac{1}{n}\sum_{j,k}(L^n)_{jk}, n \geq 1. \tag{7}$$

The first few members of the conserved operators are

$$\begin{aligned} I_1 &= \sum_j p_j, \\ I_2 &= \frac{1}{2}\sum_{j=1}^{N} p_j^2 + \sum_{1\leq j<k\leq N} \frac{a^2-a}{(x_j - x_k)^2} = H, \\ I_3 &= \frac{1}{3}\sum_j p_j^3 + \frac{1}{2}\sum_{j\neq k} \frac{a^2-a}{(x_j - x_k)^2}(p_j + p_k) \\ &- \frac{\mathbf{i}a^3}{3} \sum_{\substack{j\neq l, l\neq s \\ s\neq j}} \frac{1}{(x_j - x_l)(x_l - x_s)(x_s - x_j)}. \end{aligned} \tag{8}$$

3. Virasoro-Witt 3-algebra

The operator Nambu 3-bracket was proposed by Nambu [36],

$$\begin{aligned} [\hat{A}, \hat{B}, \hat{C}] &= \hat{A}\hat{B}\hat{C} - \hat{B}\hat{A}\hat{C} + \hat{C}\hat{A}\hat{B} - \hat{A}\hat{C}\hat{B} + \hat{B}\hat{C}\hat{A} - \hat{C}\hat{B}\hat{A} \\ &= [\hat{B}, \hat{C}]\hat{A} + [\hat{C}, \hat{A}]\hat{B} + [\hat{A}, \hat{B}]\hat{C}, \end{aligned} \tag{9}$$

where $[\hat{A}, \hat{B}] = \hat{A}\hat{B} - \hat{B}\hat{A}$. It should be mentioned that there is another definition of an operator 3-bracket [37], which is valid for operators of trace class. In this paper we only focus on the operator Nambu 3-bracket (9).

Let us introduce $N \times N$ diagonal matrix X with the coordinates x_j,

$$X = \sum_{j=1}^{N} x_j E. \tag{10}$$

where E is the unit matrix. Note that the operator matrix X is indeed the boost operator [21]. It is easy to verify that the operator matrix X and the Lax operator L satisfy the commutation relation

$$[L, X] = -\mathbf{i}E. \tag{11}$$

We define the new matrices $L^{\pm} = L \pm X$. By means of (11), we have

$$[L^+, L^-] = 2\mathbf{i}E. \tag{12}$$

In order to construct the desired infinite-dimensional 3-algebra, let us first turn to consider the following operators:

$$R_m = -\frac{1}{2\mathbf{i}}(L^+)^{m+1}L^- + mN(L^+)^m, \quad T_m = (L^+)^m. \tag{13}$$

Their commutation relations are

$$[R_m, R_n] = (n-m)R_{m+n}, \quad [R_m, T_n] = nT_{m+n}, \quad [T_m, T_n] = 0. \tag{14}$$

It can be recognized as the semidirect product of the Witt algebra with an abelian current algebra.

Based on the operator Nambu 3-bracket (9) and the commutation relations (14), after a straightforward calculation we obtain

$$\begin{aligned}
&[R_m, R_n, R_k] = (N - N^2)(k-m)(m-n)(n-k)T_{m+n+k},\\
&[R_m, R_n, T_k] = (n-m)(R_{m+n+k} + (1-2N)kT_{m+n+k}),\\
&[R_m, T_n, T_k] = (n-k)T_{m+n+k},\\
&[T_m, T_n, T_k] = 0.
\end{aligned} \tag{15}$$

Let us take the scaled generators

$$\hat{R}_m = -(N - N^2)^{-\frac{1}{4}}R_m, \quad \hat{T}_m = (N - N^2)^{\frac{1}{4}}T_m, \tag{16}$$

where the particle number N is a scaling parameter. Thus we can rewrite the 3-algebra relations (15) as

$$\begin{aligned}
&[\hat{R}_m, \hat{R}_n, \hat{R}_k] = (k-m)(m-n)(k-n)\hat{T}_{m+n+k},\\
&[\hat{R}_m, \hat{R}_n, \hat{T}_k] = (m-n)(\hat{R}_{m+n+k} + \lambda k\hat{T}_{m+n+k}),\\
&[\hat{R}_m, \hat{T}_n, \hat{T}_k] = (k-n)\hat{T}_{m+n+k},\\
&[\hat{T}_m, \hat{T}_n, \hat{T}_k] = 0,
\end{aligned} \tag{17}$$

where λ is given by

$$\lambda = \frac{(2N-1)}{(N-N^2)^{\frac{1}{2}}}. \tag{18}$$

We immediately recognize that (17) is nothing but the V-W 3-algebra derived in Ref.[28]. It should be pointed out that (17) includes the operators $\hat{R}_m$ and $\hat{T}_n$ only for $m \geq 0$ and $n \geq 1$, respectively. Precisely speaking (17) is a sub-V-W 3-algebra.

It was shown that only when $\lambda = \pm 2\mathbf{i}$, the V-W 3-algebra (17) does satisfy the following FI condition [28]:

$$[A, B, [C, D, E]] = [[A, B, C], D, E] + [C, [A, B, D], E] + [C, D, [A, B, E]]. \tag{19}$$

Let us take the scaling limit $N \to \infty$ in (18), we have $\lambda = -2\mathbf{i}$. In this way, we obtain a special V-W 3-algebra which satisfies the FI condition. It indicates that there is a well-defined special V-W 3-algebraic structure for the case of the large N limit of the quantum Calogero-Moser model.

4. $W_{1+\infty}$ 3-algebra

Let us take the generators

$$W_m^r = (-1)^r e^{-mX} L^r, \quad m, r \in \mathbb{Z}, r \geq 0. \tag{20}$$

In terms of the zero-mode operators W_0^n, the conserved operators (7) of the quantum Calogero-Moser system can be expressed as

$$I_n = \frac{(-1)^n}{n} \sum_{j,k} (W_0^n)_{jk}. \tag{21}$$

The commutation relation of the operators (20) is

$$[W_m^r, W_n^s] = (\sum_{j=0}^{r} C_r^j (-\mathbf{i}n)^j - \sum_{j=0}^{s} C_s^j (-\mathbf{i}m)^j) W_{m+n}^{r+s-j}, \tag{22}$$

where $C_r^j = \frac{r(r-1)(r-2)\ldots(r-j+1)}{j!}$.

We note that (22) is nothing but the so-called $W_{1+\infty}$ algebra [35]. It was shown that this $W_{1+\infty}$ algebra is intrinsically related to the first Hamiltonian structure of the KP hierarchy [5, 6].

Substituting the generators (20) into the operator Nambu 3-bracket (9) and using the commutation relation (22), after a tedious but straightforward calculation, we obtain the $W_{1+\infty}$ 3-algebra derived in Ref.[35]

$$
\begin{aligned}
&[W_m^r, W_n^s, W_k^h] = \\
&(\sum_{j=0}^{r} C_r^j(-\mathbf{i}n)^j - \sum_{j=0}^{s} C_s^j(-\mathbf{i}m)^j) \sum_{l=0}^{r+s-j} C_{r+s-j}^l(-\mathbf{i}k)^l W_{m+n+k}^{r+s+h-j-l} \\
&+(\sum_{j=0}^{s} C_s^j(-\mathbf{i}k)^j - \sum_{j=0}^{h} C_h^j(-\mathbf{i}n)^j) \sum_{l=0}^{s+h-j} C_{s+h-j}^l(-\mathbf{i}m)^l W_{m+n+k}^{r+s+h-j-l} \\
&+(\sum_{j=0}^{h} C_h^j(-\mathbf{i}m)^j - \sum_{j=0}^{r} C_r^j(-\mathbf{i}k)^j) \sum_{l=0}^{r+h-j} C_{r+h-j}^l(-\mathbf{i}n)^l W_{m+n+k}^{r+s+h-j-l}. \qquad (23)
\end{aligned}
$$

It is known that the FI condition does not hold for this ternary algebra. It satisfies a 3-on-3-on-3 multiple bracket relation, i.e., so-called BI condition [38]

$$
\epsilon^{i_1 i_2 \cdots i_6}[[A,[B_{i_1},B_{i_2},B_{i_3}],B_{i_4}],B_{i_5},B_{i_6}] = \epsilon^{i_1 i_2 \cdots i_6}[[A,B_{i_1},B_{i_2}],[B_{i_3},B_{i_4},B_{i_5}],B_{i_6}], \qquad (24)
$$

where $i_1, \cdots, i_6$ are implicitly summed from 1 to 6.

For the Nambu 3-bracket $[W_m^r, W_n^s, W_0^0]$, we may define it to be a parametrized bracket relation $[W_m^r, W_n^s]_{W_0^0}$. This parametrized bracket relation gives rise to the $W_{1+\infty}$ algebra (22)

$$
[W_m^r, W_n^s]_{W_0^0} \equiv [W_m^r, W_n^s, W_0^0] = [W_m^r, W_n^s]. \qquad (25)
$$

For the $W_{1+\infty}$ algebra, the intrinsic connection between this infinite-dimensional algebra and the KP hierarchy has been established. Recently the relation between the $W_{1+\infty}$ 3-algebra and the the KP hierarchy has been investigated in the framework of Nambu mechanics [35]. Some intriguing properties of the KP hierarchy have been revealed. The KP hierarchy and the CM system are the two different integrable systems. It was shown that the CM system can be obtained by a symmetry constraint of the KP hierarchy [39]. Due to the interrelation between the CM system and the KP hierarchy, it is unsurprising here that there is the W_∞ 3-algebraic structure for the quantum CM model.

Let us take the scaled generators as follows:

$$
\hat{W}_m^r = \frac{1}{N^{r-\frac{1}{2}}} W_m^r. \qquad (26)
$$

By means of (23), we obtain

$$
\begin{aligned}
&[\hat{W}_m^r, \hat{W}_n^s, \hat{W}_k^h] = \mathbf{i}(h(n-m)+s(m-k)+r(k-n))\hat{W}_{m+n+k}^{r+s+h-1} \\
&+(\sum_{j=2}^{r} C_r^j(-\mathbf{i}n)^j - \sum_{j=2}^{s} C_s^j(-\mathbf{i}m)^j) \sum_{l=0}^{r+s-j} C_{r+s-j}^l(-\mathbf{i}k)^l \frac{1}{N^{j+l-1}} \hat{W}_{m+n+k}^{r+s+h-j-l} \\
&+(\sum_{j=2}^{s} C_s^j(-\mathbf{i}k)^j - \sum_{j=2}^{h} C_h^j(-\mathbf{i}n)^j) \sum_{l=0}^{s+h-j} C_{s+h-j}^l(-\mathbf{i}m)^l \frac{1}{N^{j+l-1}} \hat{W}_{m+n+k}^{r+s+h-j-l} \\
&+(\sum_{j=2}^{h} C_h^j(-\mathbf{i}m)^j - \sum_{j=2}^{r} C_r^j(-\mathbf{i}k)^j) \sum_{l=0}^{r+h-j} C_{r+h-j}^l(-\mathbf{i}n)^l \frac{1}{N^{j+l-1}} \hat{W}_{m+n+k}^{r+s+h-j-l}. \qquad (27)
\end{aligned}
$$

By applying the scaling limit $N \to \infty$, we may drive the following 3-algebra from (27):

$$[\hat{W}^r_m, \hat{W}^s_n, \hat{W}^h_k] = \mathbf{i}(h(n-m) + s(m-k) + r(k-n))\hat{W}^{r+s+h-1}_{m+n+k}. \tag{28}$$

This is the w_∞ 3-algebra which satisfies the FI condition.

5. Summary

We reinvestigated the quantum CM model and revealed the hidden symmetries. Through the use of the boost and Lax operators of this integrable system, we showed that there are two infinite dimensional 3-algebraic structures for the quantum CM model. They are the $W_{1+\infty}$ and V-W 3-algebras, respectively. We scaled all the generators by the particle number N. In the large N limit, we found that these two infinite dimensional 3-algebras reduce to the w_∞ and special V-W 3-algebras which satisfy the FI condition, respectively. Our investigation turned out that there exists a much profound interrelation between the infinite dimensional 3-algebras and the quantum CM model. It provides new insight into this quantum many-body integrable system.

Acknowledgements

The authors are grateful to Morningside Center of Chinese Academy of Sciences for providing excellent research environment and financial support to our seminar in mathematical physics. This work is partially supported by the NSF projects (11375119 and 11475116), KZ201210028032.

References

[1] J.L. Gervais and A. Neveu, Dual string spectrum in Polyakov's quantization (II). Mode separation, Nucl. Phys. B **209** (1982) 125.

[2] J.L. Gervais, Infinite family of polynomial functions of the Virasoro generators with vanishing Poisson brackets, Phys. Lett. B **160** (1985) 277.

[3] P. Mathieu, Extended classical conformal algebras and the second hamiltonian structure of Lax equations, Phys. Lett. B **208** (1988) 101.

[4] K. Yamagishi, The KP hierarchy and extended Virasoro algebras, Phys. Lett. B **205** (1988) 466.

[5] F. Yu and Y.S. Wu, Hamiltonian structure, (anti-)self-adjoint flows in the KP hierarchy and the $W_{1+\infty}$ and W_∞ algebras, Phys. Lett. B **263** (1991) 220.

[6] K. Yamagishi, A hamiltonian structure of KP hierarchy, $W_{1+\infty}$ algebra, and self-dual gravity, Phys. Lett. B **259** (1991) 436.

[7] J.M. Figueroa-O'Farrill, J. Mas and E. Ramos, Bihamiltonian structure of the KP hierarchy and the W_{KP} algebra, Phys. Lett. B **266** (1991) 298.

[8] F. Calogero, Ground state of one-dimensional N body system, J. Math. Phys. **10** (1969)2191; Solution of a three body problem in one dimension, J. Math. Phys. **10** (1969) 2197; Solution of the one-dimensional N body problems with quadratic and/or inversely quadratic pair potentials, J. Math. Phys. **12** (1971) 419.

[9] J. Moser, Three integrable Hamiltonian systems connected with isospectral deformations, Adv. Math. **16** (1975) 197.

[10] B. Sutherland, Quantum many body problem in one dimension: ground state, J. Math. Phys. **12** (1971) 246; Exact results for a quantum many body problem in one dimension, Phys. Rev. A **4** (1971) 2019; Exact results for a quantum many body problem in one dimension II, Phys. Rev. A **5** (1972) 1372.

[11] N. Kawakami, Novel hierarchy of the SU(N) electron models and edge states of fractional quantum Hall effect, Phys. Rev. Lett. **71** (1993) 275.

[12] H. Azuma, S. Iso, Explicit relation of the quantum Hall effect and the Calogero-Sutherland model, Phys. Lett. B **331** (1994) 107.

[13] A.P. Polychronakos, Quantum Hall states as matrix Chern-Simons theory, JHEP **04** (2001) 011 [hep-th/0103013].

[14] J.A. Minahan and A.P. Polychronakos, Equivalence of two-dimensional QCD and the c=1 matrix model, Phys. Lett. B**312** (1993) 155; Interacting fermion systems from two-dimensional QCD, Phys. Lett. B**326** (1994) 288.

[15] A.P. Polychronakos, Non-relativistic bosonization and fractional statistics, Nucl. Phys. B**324** (1989) 597.

[16] M.V.N. Murthy and R. Shankar, Thermodynamics of a one-dimensional ideal das with fractional exclusion statistics, Phys. Rev. Lett. **73** (1994) 3331.

[17] B.D. Simons, P.A. Lee and B.L. Altshuler, Matrix models, one-dimensional fermions, and quantum chaos, Phys. Rev. Lett. **72** (1994) 64.

[18] G.W. Gibbons and P.K. Townsend, Black holes and Calogero models, Phys. Lett. B **454** (1999) 187.

[19] E. D'Hoker and D.H. Phong, Calogero-Moser systems in SU(N) Seiberg-Witten theory, Nucl. Phys. B **513** (1998) 405; Seiberg-Witten theory and Calogero-Moser systems, Prog. Theor. Phys. Suppl. **135** (1999) 75 [hep-th/9906027].

[20] H. Verlinde, Superstrings on AdS_2 and Superconformal Matrix Quantum Mechanics, hepth/0403024.

[21] H. Ujino, M. Wadati and K. Hikami, The quantum Calogero-Moser model: algebraic structures, J. Phys. Soc. Jpn. **62** (1993) 3035.

[22] M. Wadati and H. Ujino, The Calogero model: Integrable structure and orthogonal basis, arXiv:cond-mat/9706156.

[23] K. Hikami and M. Wadati, Integrable systems with long-range interactions, W_∞ algebra, and energy spectrum, Phys. Rev. Lett. **73** (1994) 1191; Infinite symmetry of the spin systems with inverse square interactions, J. Phys. Soc. Jpn. **62** (1993) 4203.

[24] J. Bagger and N. Lambert, Modeling multiple M2's, Phys. Rev. D **75** (2007) 045020 [hep-th/0611108]; Gauge symmetry and supersymmetry of multiple M2-branes, Phys. Rev. D **77** (2008) 065008 [arXiv:0711.0955]; Comments on multiple M2-branes, JHEP **02** (2008) 105 [arXiv:0712.3738].

[25] A. Gustavsson, Algebraic structures on parallel M2-branes, Nucl. Phys. B **811** (2009) 66 [arXiv:0709.1260].

[26] T. Curtright, D. Fairlie, X. Jin, L. Mezincescu and C. Zachos, Classical and quantal ternary algebras, Phys. Lett. B **675** (2009) 387 [arXiv:0903.4889].

[27] L. Ding, X.Y. Jia, K. Wu, Z.W. Yan and W.Z. Zhao, On q-deformed Virasoro-Witt n-algebra, arXiv:1404.0464.

[28] T.L. Curtright, D.B. Fairlie and C.K. Zachos, Ternary Virasoro-Witt algebra, Phys. Lett. B **666** (2008) 386 [arXiv:0806.3515].

[29] S. Chakrabortty, A. Kumar and S. Jain, w_∞ 3-algebra, JHEP **09** (2008) 091 [arXiv:0807.0284].

[30] V.T. Filippov, n-Lie algebras, Sib. Math. J. **26** (1985) 879.

[31] L. Takhtajan, On foundation of the generalized Nambu mechanics (second version), Commun. Math. Phys. **160** (1994) 295 [hep-th/9301111].

[32] M.R. Chen, K. Wu and W.Z. Zhao, Super w_∞ 3-algebra, JHEP **09** (2011) 090 [arXiv:1107.3295].

[33] H. Lin, Kac-Moody extensions of 3-algebras and M2-branes, JHEP **07** (2008) 136 [arXiv:0805.4003].

[34] M.R. Chen, S.K. Wang, K. Wu and W.Z. Zhao, Infinite-dimensional 3-algebra and integrable system, JHEP **12** (2012) 030 [arXiv:1201.0417].

[35] M.R. Chen, S.K. Wang, X.L. Wang, K. Wu, and W.Z. Zhao, On $W_{1+\infty}$ 3-algebra and integrable system, Nucl. Phys. B **891** (2015) 655 [arXiv:1309.4627 [nlin.SI]].

[36] Y. Nambu, Generalized Hamiltonian dynamics, Phys. Rev. D **7** (1973) 2405.

[37] H. Awata, M. Li, D. Minic and T. Yoneya, On the quantization of nambu brackets, JHEP **02** (2001) 013 [arXiv:hep-th/9906248].

[38] M.R. Bremner and L.A. Peresi, Ternary analogues of Lie and Malcev algebras, Linear Algebra Appl. **414** (2006) 1.

[39] L.V. Bogdanov and B.G. Konopelchenko, Generalized KP hierarchy: Möbius symmetry, symmetry constraints and Calogero-Moser system, arXiv:solv-int/9912005.